

EXPLICIT CONSTRUCTIONS FOR GENUS 3 JACOBIANS

JESUS ROMERO-VALENCIA & ALEXIS G. ZAMORA

ABSTRACT. Given a canonical genus three curve $X = \{F = 0\}$, we construct, emulating Mumford discussion for hyperelliptic curves, a set of equations for an affine open subset of the jacobian JX . We give explicit algorithms describing the law group in JX . Finally we introduce a related construction by means of an imbedding of the open set previously described in a Grassmanian variety.

1. INTRODUCTION

The aim of this paper is to construct, in the spirit of [7], a set of explicit equations for an affine open set of the Jacobian Variety of a canonical genus three curve X , defined by a quartic equation $F = 0$. We develop in details the case when X has a hyper-flex point, but the construction is valid for any genus 3 non-hyperelliptic curve. By a hyper-flex we understand a point $p \in X$ such that its tangent line intersects X of order 4 at p . Our ground field is $\mathbb{C}$, but the results seem to be valid over any algebraically closed field of characteristic different from 2 or 3.

The idea is to work with a subset of degree 3 non-special effective divisors and associate to it a pencil of conics. Not only an open set of JX can be described, but also the multiplication by -1 and the law of group (even when this last not explicitly in terms of affine coordinates).

In section 3 we construct an affine variety $Z \subset \mathbb{P}^{17}$ that corresponds bijectively with a subset of $Pic^3(X)$. Z is described by means of the possible decompositions $F = AG + BH$ for certain class of conics A and B . We prove that this correspondence is in fact an isomorphism of algebraic varieties.

The fact that points in an open set of the Jacobian can be put in correspondence with a decomposition of the type $F = AG + BH$ is by no means surprisingly, this follows from a homological construction, namely the computation of a minimal resolution of the $\mathcal{O}_{\mathbb{P}^2}$ -module $\mathcal{O}_X(D - 3\infty)$ for D a general element of $Pic^3(X)$ (and thus $[D - 3\infty]$ a general element of $JacX$). The degeneracy locus of this resolution is, by one hand the curve X itself, and by the other the determinant of a 2×2 matrix with coefficients in $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2))$. This interpretation is developed in section 2.

In section 4 we work with a curve with a hyper-flex point and we describe multiplication by (-1) in JX through a simple geometric construction, prove that Z is invariant under multiplication by (-1) and write explicitly the morphism:

$$Z \xrightarrow{(-1)} Z.$$

1991 *Mathematics Subject Classification.* 14H40, 14H45.

The authors were partially supported by CONACyT Grant 25811

This description allows us to obtain equations for the open set $Z/\{\pm 1\}$ of the Kummer variety $K(X)$. In that section we identify JX with the set of equivalence class of divisor of the form $D - 3\infty$, with any D effective divisors of degree 3 and ∞ a fixed hyper-flex point of X . We finish the section by describing geometrically how to obtain the sum of two divisors $D - 3\infty, D' - 3\infty \in JX$.

Section 4 is devoted to outline a more intrinsic variant of the previous construction. The idea is, instead of looking at the decomposition $F = AG + BH$, to go directly to the pencil of conics generated by A and B and define a morphism

$$Z \rightarrow \mathbb{G}(1, V),$$

with $V \subset H^0(\mathbb{P}^2, \mathcal{O}(2))$ the space of conics passing through ∞ .

Other explicit constructions of Jacobians can be found in [7], [1], [2], [4], [9]. Undoubtedly, the beautiful construction in [7] (that have its roots, according to Mumford, in works by Jacobi) have been the source of inspiration for the rest of the mentioned papers.

Even when generalizations of this construction for higher genus could be more complicated (for instance, quotients by the action of groups of matrices having as coefficients homogeneous polynomials can be involved), it is clear that the homological method of minimal resolution (see section 2) is the natural mechanism for treat such possible generalization. The discussion in genus 3 is considerable simplified by the fact that the resolution gives place to an expression of F like a 2×2 determinant. The authors are deeply indebted to an anonymous referee that indicated how our original elementary construction, developed in section 3, fits within the beautiful framework of minimal resolutions.

2 THE HOMOLOGICAL CONSTRUCTION

This section is based on [6]. Another important exposition of this ideas appears in [3].

Let $\mathcal{F}$ be a coherent sheaf on $\mathbb{P}^n$, then $\mathcal{F}$ is Cohen-Macaulay and, therefore, a minimal resolution

$$0 \rightarrow L_k \xrightarrow{\alpha_k} L_{k-1} \xrightarrow{\alpha_{k-1}} \dots \rightarrow L_0 \rightarrow \mathcal{F} \rightarrow 0,$$

with $L_p = \bigoplus_q (B_{p,q}) \otimes \mathcal{O}_{\mathbb{P}^n}(-q)$ and k the codimension of $\mathcal{F}$ in $\mathbb{P}^n$, exists. Here $B_{p,q}$ is a vector space whose dimension indicates the number of summands $\mathcal{O}_{\mathbb{P}^n}(-q)$ appearing in L_p . This minimal resolution is unique up to endomorphism of the L_i and the α_i 's are given by matrices of homogeneous polynomials. If we want to compute such a resolution it is sufficient to know the dimension of the vector spaces $B_{p,q}$.

M. Green introduced an explicit way to compute this vector spaces, based on Koszul resolution and Euler's exact sequence. Although it is not the only method appearing in the literature (see [3]) we prefer here this treatment.

We are interested in a particular case: $n = 2$, and $X \subset \mathbb{P}^2$ is a degree d curve. If $\mathcal{F}$ is any coherent sheaf on $X = \{F = 0\}$ we have a minimal resolution:

$$0 \rightarrow L_1 \xrightarrow{\alpha} L_0 \rightarrow \mathcal{F} \rightarrow 0,$$

with L_1 and L_2 of the same rank. Moreover, $\det \alpha = \lambda F$, $\lambda \in \mathbb{C}$, just because the vanishing of this determinant correspond to the geometrical locus where the map

α is not surjective. In particular, the sum of the degrees of the diagonal entries of α must equals d .

The spaces $B_{1,q}$ in the resolution of a coherent sheaf on X can be computed in the following way:

Proposition 2.1. *Let $\mathcal{O}_X(D)$ be a line bundle on X , then the terms $B_{1,q}$ in the minimal resolution of $\mathcal{O}_X(D)$ as $\mathcal{O}_{\mathbb{P}^2}$ -module are:*

$$B_{1,q} \simeq \ker \left(H^1(X, \det \Omega_{\mathbb{P}^2}(1) \otimes \mathcal{O}_X(D)(q-2)) \rightarrow \wedge^2 \mathbb{C}^3 \otimes H^1(X, \mathcal{O}_X(D)(q-2)) \right),$$

where the maps appearing in the statement are given by the restriction to X of the Koszul complex.

Proof. This is just a combination of Theorem 2.1 and Proposition 2.4 in [6].

We want to make this computation explicit in the case of a non-singular quartic on $\mathbb{P}^2$, that is, a canonical genus 3 curve. We have:

Proposition 2.2. *Let $X = \{F = 0\}$ a non-singular quartic on $\mathbb{P}^2$. Let D be a zero-degree divisor on X , $D \neq 0$, then $\mathcal{O}_X(D)$ admits a minimal resolution:*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-3)^{\oplus 2} \rightarrow \mathcal{O}_{\mathbb{P}^2}(-1)^{\oplus 2} \rightarrow \mathcal{O}_X(D) \rightarrow 0.$$

Proof. First, we compute the dimension of the vector space $B_{1,3}$. By the previous proposition it is the kernel of a map:

$$H^1(X, \det \Omega_{\mathbb{P}^2}(1) \otimes \mathcal{O}_X(D)(1)) \rightarrow \wedge^2 \mathbb{C}^3 \otimes H^1(X, \mathcal{O}_X(D)(1)).$$

Now, $\mathcal{O}_X(1) \simeq \omega_X$, thus being D not equivalent to zero we have

$$h^1(X, \mathcal{O}_X(D)(1)) = h^0(X, \mathcal{O}_X(-D)) = 0.$$

Thus, the dimension of the kernel equals the dimension of the vector space

$$H^1(X, \det \Omega_{\mathbb{P}^2}(1) \otimes \mathcal{O}_X(D)(1)).$$

Euler's exact sequence implies that $\det \Omega_{\mathbb{P}^2}(1) = \mathcal{O}_{\mathbb{P}^2}(-1)$. Therefore,

$$H^1(X, \det \Omega_{\mathbb{P}^2}(1) \otimes \mathcal{O}_X(D)(1)) \simeq H^1(X, \mathcal{O}_X(D)),$$

and, using again that D is not the zero divisor we have that $h^1(X, \mathcal{O}_X(D)) = 2$. We conclude that $\dim B_{1,3} = 2$.

Next, note that $h^1(X, \det \Omega_{\mathbb{P}^2}(1) \otimes \mathcal{O}_X(D)(k)) = h^1(X, \mathcal{O}_X(D) \otimes \omega_X^{k-1}) = 0$ for $k \geq 2$. Thus, no terms $B_{1,q}$ appears in the minimal resolution for $q \geq 4$.

Finally, for terms $B_{1,q}$ with $q < 3$ we note that the kernels to be computed are those in co-homology associated to the restricted Euler's sequence:

$$0 \rightarrow \Omega_{\mathbb{P}^2}^1(1)(D) \rightarrow \wedge^2 \mathbb{C}^3 \otimes \mathcal{O}_X(D) \rightarrow \omega_X(D) \rightarrow 0,$$

after tensor product by ω_X^{-k} , $k > 0$.

The injectivity of this maps at the 1-level co-homology follows from the fact that $h^0(X, \omega_X^{-k}(D)) = 0$ for $k \leq 0$.

Therefore $L_1 = \mathcal{O}_{\mathbb{P}^2}(-3)^{\oplus 2}$, it follows from the fact that the sum of the degree of the diagonal entries of α must be 4 that $L_0 = \mathcal{O}_{\mathbb{P}^2}(-1)^{\oplus 2}$ $\square$

In this way $\det \alpha = 0$ gives rise to a decomposition $F = AG - BH$ for some conics (A, B, G, H) . This expression is uniquely determined by D up to the actions by endomorphisms of $GL(2)$ on $\mathcal{O}_{\mathbb{P}^2}(-3)^{\oplus 2}$ and $\mathcal{O}_{\mathbb{P}^2}(-1)^{\oplus 2}$ respectively. In conclusion:

Lemma 2.2. *With the same notation as before there is a one to one correspondence between non-trivial degree-zero line bundles on X and conics (A, B, G, H) satisfying $F = AG - BH$, modulo the action of $GL(2)^{\oplus 2}$ given by:*

$$(A_0, B_0) \cdot \begin{pmatrix} A & B \\ H & G \end{pmatrix} = A_0^{-1} \begin{pmatrix} A & B \\ H & G \end{pmatrix} B_0.$$

The goal of the next section is to prove that this correspondence, at least when restricted to some open set of $JacX$, is in fact a isomorphism of algebraic varieties, and to give an explicit description of the quotient of quadruples of conics modulo the action of $GL(2)^{\oplus 2}$.

Of course, we can construct a similar correspondence at least in the case of non-singular plane curves X and obtain a set of equations that, up to the action of some algebraic group, describe, at least birationally, an open set of $JacX$. However, the explicit computation of the quotient and the resulting decomposition for the equation of X could have in general not a so simple description. For instance, if $d = 5, g = 6$ the matrix α is a 5×5 matrix of linear coefficients.

3. ALGEBRAIC EXPLICIT CONSTRUCTION FOR AN AFFINE SET OF A GENUS 3 JACOBIAN

Let $X = \{F = 0\}$ be a plane genus 3 curve defined by the non-singular homogeneous quartic polynomial F . We assume, in order to simplify the discussion, that X has a hyper-flex point, by the end of the section we shall indicate how this hypothesis can be removed. Fix, once and for all, a set of homogeneous coordinates in $\mathbb{P}^2$, in such a way that the only point of X at the infinity line $z = 0$ is $(0 : 1 : 0)$, this point will be denote by ∞ . Thus, ∞ is a hyper-flex point: $X \cdot \{z = 0\} = 4\infty$.

Denote by $Div^{3,+}(X)$ the set of degree 3 effective divisors on X , and define the following subset:

$$Div_0^{3,+}(X) := \{D = p_1 + p_2 + p_3 \in Div^{3,+}(X) \mid h^1(X, \mathcal{O}(D)) = 0, \text{ and } \infty \neq p_i\} \\ \cap \{D \in Div^{3,+}(X) \mid p_i \neq \infty \text{ and } h^1(X, \mathcal{O}(p_i + p_j + \infty)) = 0 \quad \forall i \neq j\}.$$

Note that, as the canonical divisor K_X is cut out by the linear system of lines in $\mathbb{P}^2$, the above conditions say geometrically that ∞ is not in the support of our divisors and neither, the three points in the support, nor two points and ∞ are collinear.

Moreover, the condition $h^1(X, \mathcal{O}(D)) = 0$ allows us to identify $Div_0^{3,+}(X)$ with a subset of $Pic^3(X)$. Indeed, by Riemann-Roch any divisor $D \in Div_0^{3,+}(X)$ satisfy $h^0(X, \mathcal{O}(D)) = 1$.

The next Theorem explains the basic construction of this section: we associate to any $D \in Div_0^{3,+}(X)$ the pencil of conics cutting X out in the divisor $D + \infty$. Our choice of coordinates gives an explicit form for a basis of this pencil. The reader must compare this claim with the discussion at the end of section 2.

Theorem 3.1. *There exists bijection between the sets $\text{Div}_0^{3,+}(X)$ and*

$$Z = \{(A, B, G, H) \in (H^0(\mathcal{O}_{\mathbb{P}^2}(2)))^{\oplus 4} \mid F = AG + BH\},$$

with A and B of the following particular form in affine coordinates:

$$A = a_{00} + a_{10}x + a_{01}y - x^2,$$

$$B = b_{00} + b_{10}x + b_{01}y - xy,$$

and G satisfying that its coefficient g_{11} corresponding to the monomial xy equals 1.

Proof. Given $D \in \text{Div}_0^{3,+}(X)$, we construct the pair of conics A, B considering the pencil of conics C satisfying $C.X \geq D + \infty$. In this pencil we fix a basis, namely, that formed by the only conic A in the system having as tangent line at ∞ the line $\{z = 0\}$ and the conic B in the system defined by the condition $B(1 : 0 : 0) = 0$.

Sometimes, in order to emphasize the dependence of A and B on D we write A_D, B_D instead of A and B .

This conics can be constructed in an explicit way. For instance, if $p_i \neq p_j$, for $i \neq j$ write $p_i = (x_i, y_i, 1)$ (recall that $p_i \neq \infty$). Consider the matrix:

$$M_D = \begin{pmatrix} 1 & x_1 & y_1 \\ 1 & x_2 & y_2 \\ 1 & x_3 & y_3 \end{pmatrix}.$$

This matrix is invertible, because $h^1(X, \mathcal{O}(D)) = 0$. Thus, the systems:

$$M_D \begin{pmatrix} a_{00} \\ a_{10} \\ a_{01} \end{pmatrix} = \begin{pmatrix} x_1^2 \\ x_2^2 \\ x_3^2 \end{pmatrix},$$

and

$$M_D \begin{pmatrix} b_{00} \\ b_{10} \\ b_{01} \end{pmatrix} = \begin{pmatrix} x_1 y_1 \\ x_2 y_2 \\ x_3 y_3 \end{pmatrix},$$

have unique solutions. The conics having as coefficients the previous solutions are precisely A and B .

If, for example, $D = p_1 + 2p_2$ we need to modify the system to be solved. In this case A is defined by the solutions of:

$$\begin{pmatrix} 1 & x_1 & y_1 \\ 1 & x_2 & y_2 \\ 0 & F_y(p_2) & -F_x(p_2) \end{pmatrix} \begin{pmatrix} a_{00} \\ a_{10} \\ a_{01} \end{pmatrix} = \begin{pmatrix} x_1^2 \\ x_2^2 \\ -2x_2 F_y(p_2) \end{pmatrix}.$$

Made this construction, we have the following properties:

Lemma 3.2. *For any $D \in \text{Div}_0^{3,+}(X)$, the conics A and B described above satisfy:*

- a) A is irreducible
- b) $a_{01} \neq 0$
- c) $A.B = D + \infty$
- d) *There exists $G, H \in H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2))$ such that $F = AG + BH$.*

Proof. a) The equation of A in $\mathcal{O}_{\mathbb{P}^2, \infty}$ is given by:

$$A(x, z) = a_{00}z^2 + a_{10}xz - x^2 + a_{01}z.$$

Assume A is reducible:

$$a_{00}z^2 + a_{10}xz - x^2 + a_{01}z = (a + bx + cz)(d + ex + fz),$$

comparing both expressions, we obtain:

$$\begin{aligned} ad &= 0 \\ ae + bd &= 0 \\ be &= -1. \end{aligned}$$

If $a = 0$ then $bd = 0$, but $b = 0$ is impossible, because of the third equation, thus $d = 0$. In this way:

$$A = (bx + cz)(ex + fz).$$

As $A.X \geq D$ and the two lines in the factorization contain ∞ the condition $h^1(X, \mathcal{O}(p_i + p_j + \infty)) = 0$ would be contradicted.

b) $a_{01} = 0$ implies A is reducible.

c) Just by construction.

d) Using Noether's Fundamental Theorem ([5]), it is sufficient to prove that

$$F \in \langle A, B \rangle \mathcal{O}_p,$$

for all $p \in X$. Being X and $A = 0$ non-singular, the only non trivial case to be checked is that of divisors of the form $D = p_1 + 2p_2$. In this case,

$$I_{p_i}(F, A) \geq 2 = I_{p_i}(A, B),$$

and we are done by a well known criterion ([5], Prop. 1, page 121). $\square$

The conics H and G are not uniquely determined by A and B , however, if we consider the Koszul complex:

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2} \xrightarrow{(A, B)} \mathcal{O}_{\mathbb{P}^2}(2) \oplus \mathcal{O}_{\mathbb{P}^2}(2) \rightarrow \mathcal{O}_{\mathbb{P}^2}(4) \rightarrow 0,$$

which is exact and induces an exact sequence in global sections, we see that all the possible G and H in the previous relation belong to pencils:

$$G + \lambda B \text{ and } H - \lambda A.$$

Thus, if we fix the unique value of λ such that the conic $G + \lambda B$ have coefficient $g_{11} = 1$, then we have defined uniquely the conics G and H in the relation $F = AG + BH$.

This complete the first part of the proof, namely the assignation to each $D \in \text{Div}_0^{+,3}$ of the four conics (A, B, G, H) .

Now, we make the converse construction. Assume the conics:

$$A = a_{00} + a_{10}x + a_{01}y - x^2,$$

$$B = b_{00} + b_{10}x + b_{01}y - xy,$$

satisfy the equation

$$F = AG + BH$$

for some conics G and H . We shall prove that if $A.B = D + \infty$ then $D \in \text{Div}_0^{3,+}(X)$.

Note first that $\infty \notin \text{Supp}(D)$, because the intersection index $I_\infty(A, B) = 1$. This can be proved by considering the complete ring $\hat{\mathcal{O}}_{\mathbb{P}^2, \infty} / \langle B \rangle$. Looking at the equation defining B we see that, in this algebra, the class of x is:

$$\bar{x} = \frac{b_{00}\bar{z}^2 + b_{01}\bar{z}}{1 - b_{10}\bar{z}} = (1 + b_{10}\bar{z} + b_{10}^2 + \dots)(b_{00}\bar{z}^2 + b_{01}\bar{z}).$$

Thus, $\bar{A} \in \hat{\mathcal{O}}_{\mathbb{P}^2, \infty} / \langle B \rangle$ can be expressed like:

$$\bar{A}(\bar{z}) = a_{01}\bar{z} + h(\bar{z})\bar{z}^2.$$

This prove that

$$I_\infty(A, B) = \dim_{\mathbb{C}} \hat{\mathcal{O}}_\infty / \langle A, B \rangle = 1.$$

Next, we must show that $h^1(X, \mathcal{O}(D)) = 0$. If it is not the case, then there exists a line L such that $X.L \geq D$, but this implies $A.L \geq D$ and $B.L \geq D$, it follows that A and B have a common factor L , contradicting the irreducibility of F .

A similar argument implies that $h^1(X, \mathcal{O}(p_i + p_j + \infty)) = 0$.

This two constructions are clearly one the inverse of the another. This conclude the proof of the theorem. $\square$

The next step in our construction is to prove that the set of conics previously described has a natural structure of smooth affine variety, isomorphic with an open set of $\text{Jac}X$.

Theorem 3.3. *The set Z is a smooth affine variety of dimension 3, isomorphic to a Zariski open set of $\text{Jac}X$.*

Proof. Step 1. The structure of affine variety.

The structure of affine algebraic set is given exactly by the relationship:

$$F = AG + BH.$$

More explicitly, Z is by definition a subset $Z \subset \mathbb{A}^3 \times \mathbb{A}^3 \times \mathbb{A}^5 \times \mathbb{A}^6$, with the coefficients of (A, B, G, H) corresponding to the coordinates

$$((a_{00}, a_{10}, a_{01}), (b_{00}, b_{10}, b_{01}), (g_{00}, \dots)(h_{00}, \dots)).$$

In the set of coordinates corresponding to G there appears only 5 entries, because $g_{11} = 1$. This set is defined by the equations

$$F = AG + BH,$$

and being F a fixed curve, this determines a set of 14 algebraic equations of degree two in $(a_{ij}, b_{kl}, g_{st}, h_{uv})$.

This set can be viewed, more accurately, like the image in $\mathbb{A}^{17}$ of the map:

$$\phi : \text{Div}_0^{3,+}(X) \rightarrow Z \subset \mathbb{A}^{17},$$

$$D \rightarrow (A_D, B_D, G_D, H_D).$$

Now, note that two divisors $D, D' \in \text{Div}_0^{3,+}(X)$ are linearly equivalent if and only if they are equal, which follows from the vanishing of their h^1 . In other words, $\text{Div}_0^{3,+}(X) \subset \text{Pic}^3(X)$. $\text{Div}_0^{3,+}(X)$ is in fact an open Zariski set, as can be proved by noting that in the expression:

$$\begin{aligned} & \text{Pic}^3(X) - \text{Div}_0^{3,+}(X) \\ &= \{D = p_1 + p_2 + p_3 \in \text{Div}_0^{3,+}(X) \mid h^1(X, \mathcal{O}(D)) = 1, \text{ or } \infty = p_i\} \\ & \cup \{D \in \text{Div}_0^{3,+}(X) \mid h^1(X, \mathcal{O}(p_i + p_j + \infty)) = 1 \text{ for some } i \neq j \text{ or } p_i = \infty\}, \end{aligned}$$

the first set is just the image of $W_2 \subset \text{Pic}^2(X)$ under the map $+\infty : \text{Pic}^2(X) \rightarrow \text{Pic}^3(X)$, and the second can be obtained as the epimorphic image of the morphism:

$$\pi : X \times X \rightarrow \text{Pic}^3(X),$$

defined as follows: given (p, q) , be L the line determined by q and ∞ , and set $p_2 + p_3 = X.L - q - \infty$. Then, define $\pi(p, q)$ as the class of $p + p_2 + p_3$.

Moreover, ϕ is easily expressed as a function of the symmetric functions on x_i and y_i . In this way, ϕ is an injective algebraic map onto Z . In particular, Z is irreducible.

Step 2. Dimension of Z .

Let us to compute the Zariski dimension of Z at any point and to prove it is equal to 3. Note that Z is defined by 14 equations in $\mathbb{A}^{17}$.

Given $(A, B, G, H) \in Z$, consider the tangent space:

$$\begin{aligned} T_{(A,B,G,H)}Z &= \{(\dot{A}, \dot{B}, \dot{G}, \dot{H}) \mid \deg \dot{A}, \dot{B}, \dot{G}, \dot{H} \leq 2 \text{ and} \\ & F = (A + \epsilon \dot{A})(G + \epsilon \dot{G}) + (B + \epsilon \dot{B})(H + \epsilon \dot{H}) \pmod{\epsilon^2}\} \end{aligned}$$

Thus, we must have:

$$(2.1) \quad A\dot{G} + \dot{A}G + B\dot{H} + \dot{B}H = 0.$$

Hence, our problem reduces to prove that any quartic in $\mathbb{P}^2$ containing the point $(0 : 1 : 0)$ can be expressed in the form 2.1, for some $\dot{A}, \dot{B}, \dot{G}, \dot{H}$.

Let $V = \{A\dot{G} + \dot{A}G + B\dot{H} + \dot{B}H\}$ and write

$$V = V_1 \oplus V_2,$$

with $V_1 = \{\dot{A} = \dot{B} = \dot{G} = 0\}$ and $V_2 = \{\dot{H} = 0\}$.

The space V_1 equals the space of quartics

$$\{Q \mid B \text{ divides } Q\} \simeq H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2)),$$

hence, $\dim V_1 = 6$.

Moreover, we have that inside V_2 the subspaces $V_2^0 \subset V_2$, defined by $V_2^0 = \{\dot{A} = \dot{B} = 0\}$ also have dimension 6. Thus, we only need to prove that, inside V_2 , there exists a complementary 2-dimensional subspace to V_2^0 .

It is easily provided by the space generated by H and G , that will be complementary to V_2^0 , unless $sH + tG = A$ for some constant s and t .

If this were the case, we argue in a similar fashion using now the decomposition:

$$V = \{\dot{A} = \dot{B} = \dot{H} = 0\} \oplus \{\dot{G} = 0\},$$

and use that $sH + tG = A$ and $aG + bH = B$ implies F is reducible. $\square$

In this way, we have explicitly constructed an affine variety structure for an open set of $Pic^3(X)$.

Remarks 1. Being this construction only a concrete realization of the general discussion in section 2, it seems obvious that the hypothesis on the existence of an hyper-flex is superfluous. In fact, a similar construction can be carried out if we eliminate this hypothesis, only small modifications are needed in the definition of $Div_0^{+,3}(X)$, fixing a point having as tangent $\{z = 0\}$ and avoiding divisors containing in its support any point on $z = 0$.

2. It can be proved, in the spirit of [7], that a finite number of translations of Z give an affine covering of $Pic^3(X)$, recovering the structure of $Pic^3(X)$ as a complete variety. The divisors needed for this have support on the ramifications points of the projection of X with center at ∞ on a suitable projective line. The details can be consulted in [8].

4. EXPLICIT LAW GROUP ON JX .

Let X be a canonical curve of genus 3, assume for simplicity that X has a hyper-flex point $\infty \in X$.

In this section we describe explicitly the law group on JX . The description includes an explicit algorithm for computing the inverse of an element in JX and the sum of two elements. The computation of the inverse is particularly transparent when restricted to the open set Z defined in the previous section. To start with, consider the morphism:

$$\pi_\infty : Pic^3(X) \rightarrow JX$$

$$D \rightarrow D - 3\infty.$$

It is an elementary consequence of Riemann-Roch Theorem (and also a part of Jacobi Inversion Theorem) that this map is surjective. Moreover, as previously noted, it is injective in $Div_0^{3,+}(X)$. By a slight abuse of notation we denote indistinctly by Z to both, the image in JX of $Div_0^{3,+}(X)$ and to the affine variety constructed in section 2 (they are, in fact, isomorphic).

Theorem 4.1. *The open set Z is invariant under multiplication by (-1) in JX . In the affine coordinates for Z this map is given by:*

$$(A, B, G, H) \rightarrow (A, H, G, B).$$

Proof. Let $D \in \text{Div}_0^{3,+}(X)$, denote by D^- the divisor such that $D + D^- - 6\infty \equiv 0$. This divisor can be constructed as follows: let A be the conic considered before, then $X.A = D + D' + 2\infty$ with $\deg D' = 3$. Let $L_\infty = \{z = 0\}$, being ∞ a hyper-flex we have:

$$X.A \equiv X.2L_\infty \equiv 8\infty,$$

thus, $D' = D^-$.

Let us prove that $D^- \in \text{Div}_0^{3,+}(X)$. A computation similar to the local argument invoked in the proof of Theorem 2.1 shows that $I_\infty(X, A) = 2$, thus $\infty \notin \text{Sup} D'$. Moreover, if $h^1(X, \mathcal{O}_X(D')) = 1$ then A must be reducible and the same is valid if $h^1(X, \mathcal{O}_X(q_i + q_j + \infty)) = 1$ with q_i, q_j points in the support of D' . This prove that Z is invariant under multiplication by -1 .

Now, the conic A_D coincides with A_{D^-} , as previously noted. Finally, as

$$F = AG_D + B_D H_D = AG_{D^-} + B_{D^-} H_{D^-},$$

we have

$$\begin{aligned} A.F &= D + D^- + 2\infty = A.B_{D^-} + A.H_{D^-} \\ &= D^- + \infty + A.H_{D^-} \end{aligned}$$

and thus

$$A.B_D = A.H_{D^-}.$$

By the uniqueness of the decomposition $F = AG + BH$, once fixed A and B , we have $B_{D^-} = H_D$.

In this way the set of conics associated to D^- is (A_D, H_D, G_D, B_D) . $\square$

We remark that a conic A satisfying $A.X \geq D + 2\infty$ exists independently of being $D \in \text{Div}_0^{3,+}(X)$ or not. Thus, an explicit and simple algorithm for computing the inverse of a divisor $D - 3\infty$ is to take the class of the divisor $A.X - D - 5\infty$.

The proof also shows that the conic H_D has coefficients $h_{20} = h_{02} = 0$. We obtain as a by-product:

Theorem 4.2. *The open set of the Kummer variety $K(X)$ given by the quotient $Z/\{\pm 1\}$ is determined by the equations:*

$$F = A.G + Q,$$

with Q a quartic expressible like a product of two conics passing through ∞ and $(1 : 0 : 0)$.

The condition on the reducibility of the quartic Q can be written down explicitly in terms of the usual procedure of composing the Segre map with a linear projection. More concretely, a quadric Q will be the product of such two conics if when expressed as

$$\sum_{ij} c_{ij} x^i y^j,$$

its coefficients satisfy:

$$c_{30} = c_{03} = c_{40} = c_{13} = c_{31} = c_{04} = 0,$$

$$\begin{aligned} c_{22} &= 1 & c_{00} &= w_{00}, & c_{10} &= w_{12} + w_{21} \\ c_{01} &= w_{20} + w_{02} & c_{11} &= w_{03}w_{30} + w_{12}w_{21} & c_{20} &= w_{21} \\ c_{02} &= w_{12} & c_{21} &= w_{13} + w_{31} & c_{12} &= w_{23} + w_{32} & c_{22} &= w_{33}, \end{aligned}$$

and w_{ij} satisfying the Segre relations $\text{rank}(w_{ij}) \leq 2$.

The addition law on JX , considered like the image of π_∞ , can be also described by a simple algorithm:

Let $D - 3\infty, D' - 3\infty \in JX$. Consider a cubic C such that $C.X = D + D' + E^- + 3\infty$.

Let A_{E^-} be a conic with $A_{E^-}.X = E^- + E + 2\infty$.

Now, $D + D' + E^- \equiv 9\infty$ and

$$E^- + E \equiv 6\infty.$$

It follows that $D + D' - 6\infty \equiv E + 3\infty$, that is

$$D + D' \equiv E - 3\infty.$$

Of course, Z can not be invariant under the addition law (it can not be a subgroup of JX for several reasons), so we can not expect to find a description of the sum in terms of the affine coordinates for Z .

5. THE DESCRIPTION VIA GRASSMANIANS

In this section we outline a related construction in terms of Grassmanians. This construction is a bite more intrinsic, as it does not involve the particular choice of A and B made before.

Let $V = H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2) \otimes \mathfrak{m}_\infty)$. The construction in section 3 assigns to any $D \in \text{Div}_0^{3,+}(X)$ a line in $\mathbb{P}V$, i.e., the pencil $\lambda A + \mu B$.

We have then an injective morphism:

$$Z \cong \text{Div}_0^{3,+}(X) \rightarrow \mathbb{G}(1, \mathbb{P}V),$$

$$(A, B, G, H) \rightarrow \lambda A + \mu B.$$

In terms of the Plücker embedding of the Grassmanian this morphism can be described explicitly. If we call

$$A = a_0 + a_1x + a_2y - x^2,$$

$$B = b_0 + b_1x + b_2y - xy,$$

then the Plücker embedding restricted to Z is the map sending (A, B) into

$$((a_1b_0 - b_1a_0, a_2b_0 - b_2a_0, -a_0, a_1b_2 - a_2b_1, -a_1, -a_2, -b_0, -b_1, -b_2, 1)).$$

This together with Plücker relations give another set of equations for Z . Note that it is obvious from the expression for the map that this is not only injective, but in fact an embedding when restricted to Z .

It is worthy to be noted that this map can be extended to any divisor $D \in \text{Pic}^3(X)$ such that $h^1(X, \mathcal{O}_X(D)) = 0$, as under this condition a pencil of conics cutting X out on $D + \infty$ is well defined. In other words, we can think of a morphism:

$$JX - \Theta \rightarrow \mathbb{G}(1, \mathbb{P}V).$$

This morphism, however, will be injective only on Z .

REFERENCES

1. G. W. Anderson, *An Explicit Algebraic Representation of the Abel Map*, Int. Math. Research Notices **11** (1997), 495–205.
2. A. Beauville, *Jacobians of spectral curves and completly integrable Hamiltonian system*, Acta Mathematica **164** (1990), 211–235.
3. F. Catanese, *Homological algebra and algebraic surfaces*, Algebraic geometry Santa Cruz 1995, Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI. **62** (1997), 3–56.
4. J. Estrada-Sarlabous, E. Reinaldo Barreiro, and A. Piñeiro Barceló, *On the Jacobian varieties of Picard curves: addition law and algebraic structure*, M. Nachrichten **208** (1999), 149–166.
5. W. Fulton, *Algebraic Curves*, Benjamin-Cunning, 1969.
6. M. Green, *Koszul cohomology and geometry*, ICTP College "Lectures on Riemann Surfaces" World Scientific Press (1989), 177–200.
7. D. Mumford, *Tata Lectures on Theta II*, Modern Birkhäuser Classics, 2007.
8. J. Romero-Valencia, *Construcción algebraica de la Jacobiana de una curva de género 3*, Ph. D. Thesis, CIMAT, 2008.
9. A. G. Zamora, *Some remarks on the Jacobian varieties of a Picard Curve*, Boletín SMM **3** (1997), 221–234.

UNIVERSIDAD DE GUERRERO
 AV. LÁZARO CÁRDENAS S/N
 CIUDAD UNIVERSITARIA, C.P. 39087
 CHILPANCINGO, GUERRERO, MÉXICO
E-mail address: jromv@ yahoo.com

U. A. MATEMÁTICAS, U.DE ZACATECAS
 CAMINO A LA BUFA Y CALZADA SOLIDARIDAD, C.P. 98000
 ZACATECAS, ZAC. MÉXICO
E-mail address: alexis@ mate.reduaz.mx